

Magnetic Oscillations of a Fractional Hall Dot

A.H. MacDonald^(a)

*Max-Planck-Institut für Festkörperforschung
D-7000 Stuttgart 80
Germany*

M.D. Johnson

*Department of Physics
University of Central
Florida
Orlando FL 32816-2385*

We show that a quantum dot in the fractional Hall regime exhibits mesoscopic magnetic oscillations with a period which is a multiple of the period for free electrons. Our calculations are performed for parabolic quantum dots with hard-core electron-electron interactions and are exact in the strong field limit for $k_B T$ smaller than the fractional Hall gap. Explicit expressions are given for the temperature dependence of the amplitude of the oscillations.

PACS numbers: 73.20.Dx, 73.20.Mf

Advances in nanofabrication technology have made it possible to realize artificial systems in which electrons are confined to a small area within a two-dimensional electron gas. There has been considerable interest in the physics of electron-electron and electron-hole interactions in these ‘quantum dot’ systems¹. Recent experiments have demonstrated the possibility of probing their properties in the regimes of the integer² and fractional³ quantum Hall effects⁴. The quantum Hall effect occurs whenever a system has a chemical potential discontinuity in the thermodynamic limit at a magnetic field dependent density. The fractional Hall gaps occur because of electron-electron interactions. Their microscopic origin is understood in some detail and that understanding is exploited here to discuss the thermodynamic properties of parabolic quantum dot systems in the fractional quantum Hall regime. Our calculations are exact within the hard-core model^{5,6} which may be considered as an ideal model of the fractional quantum Hall effect. We show that in the regime where the electronic ground state of the dot is the Laughlin⁷ state which occurs at filling factor $\nu = 1/m$ in the bulk, all thermodynamic properties of electrons in the dot have a component which is periodic in the magnetic flux through the area of the dot with period $m\Phi_0$. (Here $\Phi_0 \equiv hc/e$ is the electronic magnetic flux quantum.)

We consider a system of electrons in contact with a particle reservoir with chemical potential μ and confined to a finite area of a two dimensional electron gas by a parabolic potential, $V(r) = \frac{1}{2}m\Omega^2 r^2$. We measure the chemical potential with respect to the two-dimensional subband energy and define a nominal radius, R , for the quantum dot system in terms of the radius where the parabolic confining potential equals the amount by which the chemical potential exceeds the quantized kinetic energy of the lowest Landau level; $\frac{1}{2}m\Omega^2 R^2 \equiv \mu - \hbar\omega_c/2$. (Here $\omega_c = eB/mc$ is the cyclotron frequency.) In the strong magnetic field limit where $\Omega/\omega_c \ll 1$ only the states in the lowest Landau level are relevant. (Here $\omega_c = eB/mc$ is the cyclotron frequency.) In the symmetric gauge the single particle states ($\phi_l(z) \sim z^l \exp(-z\bar{z}/4\ell^2)$) in this level may be labeled by angular momentum and

have energy¹ $\epsilon_l = \hbar\omega_c/2 + \gamma(l+1)$ where $\gamma = m\Omega^2\ell^2$. (Here $z = x + iy$ is the 2D electron coordinate expressed as a complex number, $\ell \equiv (\hbar c/eB)^{1/2}$ is the magnetic length, and the allowed values of single particle angular momentum within the lowest Landau level are $l = 0, 1, 2, \dots$)

Many-fermion states in the lowest Landau level may be described in a boson language^{8,9} which will prove useful in the fractional Hall regime. The mapping between boson and fermion descriptions can be understood in terms of a useful property of many-fermion wavefunctions within the lowest Landau level. Any many-fermion wavefunction in this level may be expressed in the form

$$\Phi[z] = P(z_1, z_2, \dots, z_N) \prod_k \exp(-z_k \bar{z}_k / 4\ell^2) \quad (1)$$

where $P[z]$ is an antisymmetric polynomial. Since $P[z]$ must vanish whenever any two coordinates are identical it follows that

$$P[z] = \prod_{i < j \leq N} (z_i - z_j) Q(z_1, z_2, \dots, z_N) \quad (2)$$

where $Q[z]$ is a symmetric polynomial. The fractional quantum Hall effect is related to the existence of classes of low-energy many-body states where the probability amplitude to find any pair of electrons in a state of relative-angular-momentum less than m is identically zero. States in these classes have wavefunctions of the form

$$P[z] = \prod_{i < j \leq N} (z_i - z_j)^m Q(z_1, z_2, \dots, z_N) \quad (3)$$

In the bulk such states exist only^{10,8,11} when the Landau level filling factor $\nu \leq 1/m$ and the chemical potential jumps with increasing density at $\nu = 1/m$ as a consequence. In hard-core models^{6,5} electrons interact only when their relative-angular-momentum is less than $1/m$ so that the interaction energy is identically zero in the corresponding class of low-energy states. For a quantum dot with average $\nu < 1/m$ only these low-energy states

are relevant at temperatures low compared to the fractional Hall gap and it is useful to consider a model where the high-energy states are excluded from the Hilbert space. (The relationship between hard-core models and realistic models is discussed in more detail below.) The products $Q[z] \prod_k \exp(-z_k \bar{z}_k / 4\ell^2)$ are wavefunctions for N bosons occupying states with $l \geq 0$. Since the low-lying excitations are completely described in terms of the set of symmetric polynomials Q , it follows that for $\nu < 1/m$, the number of low-energy N -fermion wavefunctions at total angular momentum M_F equals the number of independent N -boson wavefunctions, $g(N, M_B)$, at total angular momentum

$$M_B = M_F - mN(N-1)/2. \quad (4)$$

As we show below many properties of an ideal parabolic quantum dot in the integer or fractional quantum Hall regimes can be expressed in terms of $g(N, M_B)$.

We evaluate $g(N, M_B)$ using an iterative procedure in which single-particle angular momenta are added incrementally to the boson Hilbert space. Define $g_k(N, M_B)$ to be the value of $g(N, M_B)$ when only single-particle angular momenta from 0 to k are included in the boson Hilbert space. $g_0(N, M) = \delta_{M,0}$ since there is only one boson state for any number of electrons and it has total angular momentum equal to zero. When an additional single particle state is added to the boson Hilbert space we have the possibility of having any number of bosons in that state:

$$g_k(N, M) = \sum_{N'=0}^N g_{k-1}(N - N', M - kN') \quad (5)$$

(Note that $g_k(N, M) \equiv 0$ for $M < 0$.) For $k \geq M$, $g_k(N, M) = g(N, M)$ is independent of k . Also, for $N \geq M$, $g_k(N, M) \equiv g_k(M)$ is independent of N . Combining these, for $k, N \geq M$, $g_k(N, M) = g(M)$. In Fig.[1] we plot $s(N, M) \equiv \ln g(N, M)$ against $M^{1/2}$ for several values of N . For $N \geq M$, the total number of states of boson angular momentum M can also be identified from the coefficients of the Taylor series expansion for the boson

partition function¹²

$$Z_k^B(\beta\gamma) = \prod_{k'=1}^k \frac{1}{1 - \exp(-\beta\gamma k')} = \sum_{M=0}^{\infty} g_k(M) \exp(-\beta\gamma M). \quad (6)$$

As shown in Fig.[1], for large M $s(M) \sim \sqrt{2/3\pi} M^{1/2} \sim 2.5650997 M^{1/2}$ with logarithmic corrections. This result can be obtained by comparing the mean values of the total boson momentum calculated from the two forms for the right-hand side of Eq. (6) in the limit of large k and high temperatures. In this limit, $\ln Z^B \rightarrow \pi^2/(6\beta\gamma)$.

To evaluate thermodynamic properties of the quantum dot we need only note that the energy of a fermion state in the class specified by Eq. (3) is determined by the total number of electrons and the total angular momentum:

$$E(N, M_F) = \gamma(N + M_F) = \gamma[N + mN(N-1)/2 + M_B]. \quad (7)$$

It follows that the grand partition function for the dot is given by

$$Z_G = \sum_{N=0}^{\infty} \sum_{M_B=0}^{\infty} P(N, M_B), \quad (8)$$

where

$$P(N, M_B) = g(N, M_B) \exp(\beta\gamma[NN_\phi - mN(N-1)/2 - N]) \exp(-\beta\gamma M_B), \quad (9)$$

$\beta = 1/(k_B T)$, and

$$N_\phi = (\mu - \hbar\omega_c/2)/\gamma = B\pi R^2/\Phi_0. \quad (10)$$

N_ϕ is the number of flux quanta passing through the nominal area of the dot. Thermodynamic mean values of any quantity f expressible in terms of M_F or N are now easily evaluated:

$$\langle f(N, M_F) \rangle = Z_G^{-1} \sum_{N=0}^{\infty} \sum_{M_B=0}^{\infty} f[N, M_B + mN(N-1)/2] P(N, M_B). \quad (11)$$

In Fig.[2] we show plots of the N_ϕ dependence¹³ of the magnetization of the dot for $m = 1$. We remark that in this system the magnetization $M = \langle E \rangle / B$.¹⁴ Note that as expected the magnetization has a component which is periodic in N_ϕ with period one. The results for $m = 1$ are closely related to those for non-interacting electrons discussed previously by Sivan and Imry¹⁵ and are precisely equivalent to those trivially obtained by treating the system as a free fermion system. The equivalence of the present Bose description and the Fermi description is closely related to the bosonization transformations¹⁶ which permit the exact treatment of models of a one-dimensional electron gas. (See below.) The utility of the Bose description is in treating the case $m \neq 1$, where the system is *not* equivalent to a free Fermi gas. In Fig.[3] the mesoscopic magnetization oscillations are shown for the case of a fractional Hall quantum dot with $m = 3$. Note that in this case the magnetization has a component periodic in N_ϕ with period three.

We can derive an analytic expression for the magnetization which is valid for $N_\phi \gg 1$. In this limit $g(N, M_B) = g(M_B)$ for all M_B contributing importantly to the partition function. Then the grand partition function factors into a boson part which describes all neutral excitations of the system and a part which describes the charged excitations:

$$Z_G = Z_B \exp(-\beta\Omega^*) \sum_{N=0}^{\infty} f(N + 1/2), \quad (11)$$

where $f(x) = \exp[-\beta\gamma m(x - x^*)^2/2]$, $x^* = N_\phi/m + 1 - 1/m$, and $\Omega^* = -\gamma[N_\phi^2 - 2N_\phi(1 - m/2) + (1 - m/2)^2]/2m$. The boson excitations represent charge density oscillations at the edge of the dot^{16,17}. The sum over particle numbers can be performed using the Poisson summation formula with the result

$$Z_G = Z_B \exp(-\beta\Omega^*) \left(\frac{2\pi}{\beta\gamma m} \right)^{1/2} \left[1 + 2 \sum_{s=1}^{\infty} (-)^s \exp(-2\pi^2 s^2 / \beta\gamma m) \cos(2\pi(N_\phi - 1)s/m) \right]. \quad (12)$$

Except at very low temperatures the first term inside square brackets will dominate. An

elementary calculation then leads to the following expression for the magnetization:

$$M = \frac{\gamma}{B} \left[\frac{\pi^2}{6(\beta\gamma)^2} + \frac{N_\phi^2}{2m} - \frac{(1 - m/2)^2}{2m} + \frac{1}{2\beta\gamma} \right] + M_{osc} \quad (13)$$

where

$$M_{osc} = \frac{\gamma}{B} \left[\frac{4\pi N_\phi}{m\beta\gamma} \sin(2\pi(N_\phi - 1)/m) + \frac{4\pi^2}{m(\beta\gamma)^2} \cos(2\pi(N_\phi - 1)/m) \right] \exp(-2\pi^2/\beta\gamma m) \quad (14)$$

This expression for the magnetization agrees well with the numerical results shown in Fig.[2] and Fig.[3].

For a given chemical potential and parabolic confinement the ground state is determined by a competition between the confinement energy, which favors ground states of higher density (low total angular momentum) and the interaction energy which favors ground states of lower density (higher total angular momentum.) This competition has been discussed on the basis of exact diagonalization studies for extremely small quantum dots by Maksym and Chakraborty¹⁸. For fixed chemical potential, the area of the dot $2\pi(\mu - \hbar\omega_c/2)/(m\Omega^2)$ will decrease with magnetic field, hence decreasing the strength of the confinement. The ground state for fixed electron number will jump between different incompressible states. For the hard-core model the ground state will be the $m = 3$ Laughlin state whenever the confinement potential at the edge of the dot is weak compared to the hard core potential strength.¹⁹ For realistic Coulomb interactions the dot density is determined primarily, except at extremely small dot sizes, by a competition between the Hartree interactions in the dot and the confining potential. The average filling factor in the dot at strong fields will be approximately the lesser of one and¹⁹

$$\nu = \langle N \rangle^{1/3} \left(\frac{3\pi\sqrt{2}\gamma\ell}{8e^2} \right)^{2/3}. \quad (15)$$

When correlations which favor particular densities are taken into account the ground state for fixed N will be well-approximated by the finite N Laughlin wavefunctions over finite

ranges of magnetic field strength when the average filling factor is near $1/m$. We expect that the results shown in Fig.[3] will apply for realistic interactions when the average filling factor is near $1/3$. Experimentally determined values² of the curvature of the parabolic potentials give values of γ/k_B in excess of $0.1K$ so that the effects shown in Fig.[2] and Fig.[3] should be visible at accessible temperatures. Similar mesoscopic oscillations occur in all thermodynamic properties¹⁹ of the quantum dots and presumably in all physical properties. For example we may expect that the conductance of the dot, which is more easily measured than thermodynamic properties of the dot, is closely related²⁰ to the particle number fluctuations in the dot.

The oscillations shown in Fig.[3] reflect a fundamental property of fractional Hall states. Near $\nu = 1/3$ the properties of the system in contact with a particle reservoir have contributions which are periodic in N_ϕ with period three. These oscillations express the fact that on average one electron is added to the system for each three additional flux quanta. Hence they can be observed only when the field dependence is studied at fixed chemical potential rather than fixed particle number. The quantized Hall conductance is proportional to the charge added to the system per flux quantum in the thermodynamic limit, *e.g.*, $e/3$ for $\nu = 1/3$. To obtain this result using an effective theory of non-interacting fermions²¹ would require the fermions to have a charge $e/\sqrt{3}$. Such a theory would give mesoscopic oscillations with an incorrect period.

This work was supported in part by the National Science Foundation under grant DMR-9113911. AHM acknowledges helpful conversations with Ian Affleck, M. A. Kastner, Daniela Pfannkuche and Xiao-Gang Wen and the hospitality of the University of New South Wales where part of this work was completed.

REFERENCES

- (a) Permanent address: Department of Physics, Indiana University, Bloomington IN, 47401.
1. For recent reviews see U. Merkt, *Advances in Solid State Physics*, **30**, 77 (1990); Tapash Chakraborty, *Comments on Condensed Matter Physics* **16**, 35 (1992); M.A. Kastner, *Rev. Mod. Phys.* **64**, 849 (1992).
 2. P.L. McEuen, E.B. Foxman, U. Meirav, M.A. Kastner, Y. Meir, Ned S. Wingreen, and S.J. Wind, *Phys. Rev. Lett.* **66**, 1926 (1991); P.L. McEuen, E.B. Foxman, Jari Kinaret, U. Meirav, M.A. Kastner, Ned S. Wingreen, and S.J. Wind, *Phys. Rev. B* **45**, 11419 (1992).
 3. W. Hansen, T.P. Smith III, K.Y. Lee, J.A. Brum, C.M. Knoedler, J.M. Hong, and D.P. Kern, *Phys. Rev. Lett.* **62**, 2168 (1989).
 4. K. von Klitzing, G. Dorda, and M. Pepper, *Phys. Rev. B* **45**, 494 (1980); D.C. Tsui, H.L. Stormer, and A.C. Gossard, *Phys. Rev. Lett.* **48**, 1559 (1982).
 5. S.A. Trugman and S.A. Kivelson, *Phys. Rev. B* **31**, 5280, (1985).
 6. F.D.M. Haldane, *The Quantum Hall Effect*, R.E. Prange and S.M. Girvin eds., (Springer Verlag, New York, 1986), Chapter 8.
 7. R.B. Laughlin, *Phys. Rev. Lett.*, **50**, 1395 (1983).
 8. A.H. MacDonald and D.B. Murray, *Phys. Rev. B* **32**, 2707 (1985).
 9. X.C. Xie, Song He, and S. Das Sarma, *Phys. Rev. Lett.* **66**, 389 (1991).
 10. See for example, A. H. MacDonald, *A Perspective on the Quantum Hall Effect*, (Jaca Books, Milano, 1989).

11. A.H. MacDonald, *Phys. Rev. Lett.* **64**, 222 (1990).
12. This is the boson grand partition function for bosons occupying states $k' = 1$ to k with the chemical potential set to the energy of the $k = 0$ state. This partition function accounts for the neutral phonon-like collective excitations of the system. For $N > M$ the neutral excitations and charged excitations of the system are independent so that the partition function may be factored for large systems or low temperatures (see text).
13. $BdN_\phi/dB = N_\phi - \hbar\omega_c/2\gamma$. In the strong magnetic field limit the number of flux quanta through a dot decreases with field because kinetic energy quantization depopulates the dot.
14. We give the magnetization and energy with respect to the values for free electrons in the lowest Landau level, $M_{free} = -\mu_B N$.
15. U. Sivan and Y. Imry, *Phys. Rev. Lett.* **61**, 1001 (1988).
16. F.D.M. Haldane, *J. Phys. C* **14**, 1585 (1981).
17. Michael Stone, *Phys. Rev. B* **42**, 8299 (1990); *Ann. Phys. (NY)* **207**, 38 (1991);
Michael Stone, H.W. Wyld, and R.L. Schult, *Phys. Rev. B* **45**, 14156 (1992).
18. P. A. Maksym and Tapash Chakraborty, *Phys. Rev. Lett.* **65**, 108 (1990).
19. A.H. MacDonald and M.D. Johnson, unpublished.
20. J.M. Kinaret, Y. Meir, N.S. Wingreen, P. Lee, and X.G. Wen, *Phys. Rev. B* **45**, 9489 (1992).
21. X.G. Wen, review article submitted to *Int. J. Mod. Phys. B*, and references therein.

FIGURE CAPTIONS

1. $s(N, M) \equiv \ln g(N, M)$ vs. $M^{1/2}$. The larger ‘entropy’ is for $N = \infty$ and the smaller is for $N = 10$. The dashed line is $\sqrt{2/3\pi}M^{1/2}$.
2. Magnetic oscillations of non-interacting electrons in a parabolic quantum dot at $\beta\gamma = 10$ and at $\beta\gamma = 1$. (The curve with larger oscillations is for lower temperature.) The plotted quantity is $M + \mu_B N - (\gamma/2B)(N_\phi^2 - 1/4)$. These results are valid for $\beta\hbar\omega_c \gg 1$.
3. Magnetic oscillations of electrons with hard-core interactions in a parabolic quantum dot with $\nu < 1/3$ at $\beta\gamma = 10$ and at $\beta\gamma = 1$. The plotted quantity is $M + \mu_B N - (\gamma/6B)(N_\phi^2 - 1/4)$. These results are valid for $\beta V \gg 1$ where V is the strength of the hard core interaction.